

A service of the U.S. National Library of Medicine  
and the National Institutes of HealthMy NCBI  
[Sign In] [Register]

## Exhibit D

All Databases	PubMed	Nucleotide	Protein	Genome	Structure	OMIM	PMC	Journals	Books
---------------	--------	------------	---------	--------	-----------	------	-----	----------	-------

Search  for

Advanced Search

Display   20

All: 1 Review: 0

1: Exp Mol Pathol. 1998 Oct;65(2):53-63.

Full Text Links

**Despite its homology to angiotensin apolipoprotein(a) does not affect angiogenesis.**

Lou XJ, Kwan HH, Prionas SD, Yang ZJ, Lawn RM, Fajardo LE.

Falk Cardiovascular Research Center, Stanford University School of Medicine, Stanford, California 94305, USA.

Apolipoprotein(a) [apo(a)] contains a kringle domain(IV) homologous to that of angiotensin, a natural angiogenic inhibitor. Because of this structural similarity we suspected that apo(a) could be an inhibitor of angiogenesis. The possible role of apo(a) in microvascular proliferation was studied in an in vivo quantitative model, the disc angiogenesis system (DAS) and compared to angiotensin. Apo(a) and other test compounds were placed in the center of a polyvinyl alcohol foam disc that was implanted subcutaneously in mice. After 14 days, the disc was removed and vascular growth into the disc was measured. Apo(a) did not affect spontaneous vessel growth into the disc, while angiotensin suppressed this growth and basic fibroblast growth factor (bFGF) increased it. Additionally, apo(a) did not modify the vascular growth induced by bFGF. Transgenic mice expressing the human apo(a) gene were used to study the systemic effect of apo(a): neither an increase nor a decrease in vascular growth was detected. Our results suggest that apo(a) is unlikely to play a significant role in the control of angiogenesis. Furthermore, our experiments confirm the inhibitory effect of angiotensin not only on induced angiogenesis but also on baseline, spontaneous angiogenesis. Copyright 1998 Academic Press.

PMID: 9828147 [PubMed - indexed for MEDLINE]

**Related articles**

- ▶ Inhibition of angiogenesis and angiogenesis-dependent tumor growth by [J Biol Chem. 2003] fragments of human apolipoprotein(a).
- ▶ Generation of biologically active angiotensin kringle 1-3 by activated hum [J Immunol. 2002]
- ▶ Angiogenic balance in human melanoma: expression of VEGF, bFGF [Int J Cancer. 2000] angiotensin in relation to vascular density of xenografts in vivo.
- ▶ **Review** Angiotensin: an endogenous inhibitor of angiogenesis and of tumor grow [EKS. 1997]
- ▶ **Review** Characterization and applications of the disc angiogenesis sy [Exp Mol Pathol. 1992]

» See reviews... | » See all...

**Recent Activity** 

Despite its homology to angiotensin apolipoprotein(a) does not affect angiogenesis.

Angiogenesis is impaired by hypercholesterolemia: role of asymmetric dimethylarginine.

Apolipoprotein(a), a link between atherosclerosis and tumor angiogenesis.

Secreted human apolipoprotein(a) kringle IV-10 and kringle V inhibit angiogenesis and xeno...

Lipoprotein(a) and cancer: anti-neoplastic effect besides its cardiovascular potency.

» See more...

Display   20

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)